

NATURALITY AND INDUCED REPRESENTATIONS

SIEGFRIED ECHTERHOFF, S. KALISZEWSKI, JOHN QUIGG, AND IAIN RAEBURN

ABSTRACT. We show that induction of covariant representations for C^* -dynamical systems is natural in the sense that it gives a natural transformation between certain crossed-product functors. This involves setting up suitable categories of C^* -algebras and dynamical systems, and extending the usual constructions of crossed products to define the appropriate functors. From this point of view, Green's Imprimitivity Theorem identifies the functors for which induction is a natural equivalence. Various special cases of these results have previously been obtained on an *ad hoc* basis.

1. INTRODUCTION

Induced representations and the imprimitivity theorems which characterize them are a fundamental tool in the representation theory of dynamical systems and crossed products. In the powerful formulation of Rieffel and Green, induction is done by tensoring with a Hilbert bimodule, and the imprimitivity theorem tells us how to expand the left action to make this bimodule an imprimitivity bimodule. In recent applications, it has been necessary to know that this induction process is compatible with other constructions involving crossed products. Verifying such compatibility can be painful: it is often obvious that everything must work because “induction is natural”, yet technically hard to sort out the details. And afterwards one is left feeling that one must have missed the point: the techniques are vaguely familiar even if the particular application isn't.

We have found that it is more satisfactory to phrase our questions and results directly in terms of Hilbert bimodules, rather than in terms of induced representations themselves. Thus in [KQR97] we viewed a Hilbert A – B bimodule X as an arrow from A to B , and said that a diagram

$$(1.1) \quad \begin{array}{ccc} A & \xrightarrow{X} & B \\ \downarrow Y & & \downarrow W \\ C & \xrightarrow{Z} & D \end{array}$$

Date: July 27, 1999; revised September 27, 1999.

1991 *Mathematics Subject Classification.* Primary 46L55.

This research was partially supported by the Australian Research Council, and the Office of the Vice Provost for Research at Arizona State University.

of Hilbert bimodules “commutes in the strong sense” if $Y \otimes_C Z \cong X \otimes_B W$ as A – D bimodules; if so, the Rieffel induction processes $\pi \mapsto X\text{-Ind}(W\text{-Ind } \pi)$ and $\pi \mapsto Y\text{-Ind}(Z\text{-Ind } \pi)$ yield representations which are systematically equivalent in a way compatible with intertwining maps, direct-sum decompositions and continuity.

Here we make more precise the idea that we have been verifying the naturality of induction. Category theory tells us exactly what mathematicians should mean when they talk about naturality, and how we need to set things up to make sense of it. Here we build a category in which the objects are C^* -algebras, in which the morphisms are given by Hilbert bimodules, and in which the commutativity of a diagram like (1.1) says precisely that $Y \otimes_C Z \cong X \otimes_B W$. We then apply this framework to the induction process arising from Green’s imprimitivity theorem [Gre78, Proposition 3], proving that that theorem actually gives a natural transformation between certain crossed-product functors which take values in this category. This theorem includes as special cases a number of results in the recent literature which could be paraphrased as saying “Green induction is compatible with Morita equivalence”.

Part of our point here is that taking the trouble to formulate theorems in this categorical framework will result in a theory which is more robust and more directly applicable. However, many applications require us to consider coactions or twisted crossed products, and setting up the appropriate categories, functors, and naturality theorems for these situations can take a good deal of technical effort. So in order to illustrate our approach, we concentrate here on one naturality theorem which requires only standard techniques, and leave equivariant versions involving coactions and twists to a more comprehensive sequel. We have tried hard to make this paper accessible to anyone familiar with the basic material on Hilbert modules and Morita equivalence implicit in the formulation of Green’s theorem; this can be found, for example, in the early chapters of [RW98].

We begin in Section 2 by discussing our category \mathcal{C} . The objects in \mathcal{C} will be C^* -algebras, a morphism from one C^* -algebra A to another B will be given by a Hilbert A – B bimodule X , and the composition of morphisms will be given by the balanced tensor product of bimodules. There is asymmetry here: X will be a right Hilbert B -module, and A will act by adjointable operators on X , in the sense that there is a nondegenerate homomorphism $\kappa : A \rightarrow \mathcal{L}(X_B)$ describing the action. We call such a bimodule X a right-Hilbert bimodule to stress that the Hilbert module structure is on the right.¹ The morphisms will actually be isomorphism classes $[_A X_B]$ of these right-Hilbert bimodules; this is necessary, because, for example, we expect ${}_B B_B$ to give the identity morphism on B , so ${}_A X_B$ and ${}_A (X \otimes_B B)_B$ should define the same morphism in \mathcal{C} . In Section 2 we also show that the isomorphisms in \mathcal{C} are exactly the (isomorphism classes of) imprimitivity bimodules, and prove that every morphism

¹These and related objects have been called *Hilbert bimodules* (for example, in [Pim97]), or *C^* -correspondences* (for example, in [MS98]), but we have preferred to stick with the name used in [KQR97] to stress that the Hilbert-module structure is on the right.

$[{}_AX_B]$ in \mathcal{C} can be factored as $[{}_CY_B] \circ [\varphi]$, where ${}_CY_B$ is an imprimitivity bimodule and $\varphi : A \rightarrow M(C)$ is a nondegenerate homomorphism.

In Section 3 we show how to view the crossed-product construction as a functor. We first have to add actions to the objects and morphisms of \mathcal{C} to build a category $\mathcal{A}(G)$ whose objects are dynamical systems associated with a fixed locally compact group G ; forming the crossed product then gives a functor $(A, \alpha) \mapsto A \rtimes_\alpha G$ from $\mathcal{A}(G)$ to \mathcal{C} . Because the objects in \mathcal{C} are C^* -algebras rather than isomorphism classes of C^* -algebras, we have to be careful to nominate a particular C^* -algebra as the crossed product. We discuss our nominee and its relationship to the universally defined crossed product which we usually prefer. Functoriality requires that we give a parallel construction of crossed products of Hilbert modules; the construction we have chosen is different from that of [Com84] and [CMW84] and may be of some independent interest.

Our main theorem is proved in Section 4. Green's construction associates to each object (A, α) in $\mathcal{A}(G)$ and each closed subgroup H of G a right-Hilbert $(A \rtimes_\alpha G)$ – $(A \rtimes_\alpha H)$ bimodule $X_H^G(A, \alpha)$; his Imprimitivity Theorem says that $X_H^G(A, \alpha)$ is also a $(C_0(G, A) \otimes_{\alpha \otimes \tau} G)$ – $(A \rtimes_\alpha H)$ imprimitivity bimodule. Our theorem says that the assignment $(A, \alpha) \mapsto [X_H^G(A, \alpha)]$ is a natural transformation. To prove this, we have to show that every morphism in $\mathcal{A}(G)$ gives rise to a certain commutative diagram in \mathcal{C} . We do this by factoring the morphism into an ordinary homomorphism and an imprimitivity bimodule; handling the homomorphism is straightforward, and we deal with the imprimitivity bimodule by adapting a powerful linking-algebra technique from [ER96, §4]. We find it intriguing that ideas developed to meet the demands of nonabelian duality are now feeding back into the theory of ordinary crossed products — both in the technical sense, as in our use of the linking-algebra technique, and in the motivational sense, in that we were led to formulate our theorem through our attempts to handle more complicated problems involving coactions.

2. THE CATEGORY \mathcal{C}

Let B be a C^* -algebra. A *Hilbert B -module* is a vector space X which is a right B -module equipped with a positive definite B -valued sesquilinear form $\langle \cdot, \cdot \rangle_B$ satisfying

$$(2.1) \quad \langle x, y \cdot b \rangle_B = \langle x, y \rangle_B b \quad \text{and} \quad \langle x, y \rangle_B^* = \langle y, x \rangle_B$$

for all $x, y \in X$ and $b \in B$, and which is complete in the norm $\|x\| = \|\langle x, x \rangle_B\|_B^{\frac{1}{2}}$. For the general theory of Hilbert modules, we refer to [Lan95] or [RW98].

Definition 2.1. Let A and B be C^* -algebras, and let X_B be a Hilbert module which is *full* in the sense that $\overline{\langle X, X \rangle_B} = B$. We say that X is a *right-Hilbert A – B bimodule* if it is a nondegenerate left A -module satisfying

$$(2.2) \quad a \cdot (x \cdot b) = (a \cdot x) \cdot b \quad \text{and} \quad \langle a \cdot x, y \rangle_B = \langle x, a^* \cdot y \rangle_B$$

for all $a \in A$, $x, y \in X$, and $b \in B$. (This is equivalent to having a nondegenerate homomorphism $\kappa: A \rightarrow \mathcal{L}(X_B)$ and putting $a \cdot x = \kappa(a)x$.) An *isomorphism* of right-Hilbert bimodules is a bijective linear map $\Phi: {}_A X_B \rightarrow {}_A Y_B$ such that

- (i) $\Phi(a \cdot x) = a \cdot \Phi(x)$,
- (ii) $\Phi(x \cdot b) = \Phi(x) \cdot b$, and
- (iii) $\langle \Phi(x), \Phi(y) \rangle_B = \langle x, y \rangle_B$.

(Property (ii) is redundant — it follows from (iii) by considering $\|\Phi(x \cdot b) - \Phi(x) \cdot b\|^2$ — but we include it to emphasize that Φ is a bimodule homomorphism.)

Example 2.2. If B is a C^* -algebra, then B becomes a right-Hilbert B – B bimodule by putting

$$a \cdot b \cdot c = abc \quad \text{and} \quad \langle a, b \rangle_B = a^*b$$

for $a, b, c \in B$. More generally, if $\varphi: A \rightarrow M(B)$ is a nondegenerate homomorphism, then B_B becomes a right-Hilbert A – B bimodule with left action given by

$$a \cdot b = \varphi(a)b.$$

The morphisms from A to B will be the isomorphism classes of right-Hilbert A – B bimodules (denoted with square brackets); we need to pass to isomorphism classes to show that composition of morphisms has the necessary properties. Two ordinary homomorphisms may give the same morphism in \mathcal{C} , but only if they differ by an inner automorphism:

Proposition 2.3. *Let φ and ψ be nondegenerate homomorphisms of A into $M(C)$. Then $[\varphi] = [\psi]$ in \mathcal{C} if and only if there exists $u \in UM(C)$ such that $\psi = \text{Ad } u \circ \varphi$.*

Proof. If $\psi = \text{Ad } u \circ \varphi$, then the map $c \mapsto uc$ is Hilbert module automorphism of C_C which intertwines the left actions coming from φ and ψ .

For the converse, suppose $[\varphi] = [\psi]$ in \mathcal{C} , so there exists a linear bijection $L: C \rightarrow C$ such that

$$L(cd) = L(c)d, \quad L(c)^*L(d) = c^*d, \quad \text{and} \quad L(\varphi(a)c) = \psi(a)L(c)$$

for each $c, d \in C$. Define $R: C \rightarrow C$ by $R(c) = L^{-1}(c^*)^*$. Then the first two of these properties imply that (L, R) is an invertible double centralizer of C , so there exists an invertible element u of $M(C)$ such that $L(c) = uc$ for all c . Since

$$u^{-1}c = L^{-1}(c) = L^{-1}(c^{**})^{**} = R(c^*)^* = (c^*u)^* = u^*c,$$

it follows that u is unitary, and $L(\varphi(a)c) = \psi(a)L(c)$ implies that $\psi = \text{Ad } u \circ \varphi$. \square

To define composition we use the internal tensor product of Hilbert modules. Let ${}_A X_B$ and ${}_B Y_C$ be right-Hilbert bimodules. Then the algebraic tensor product $X \odot Y$ becomes a not-necessarily-complete right-Hilbert A – C bimodule with actions given by

$$a \cdot (x \otimes y) = a \cdot x \otimes y \quad \text{and} \quad (x \otimes y) \cdot b = x \otimes y \cdot b$$

and a not-necessarily-definite C -valued inner product given by

$$\langle x \otimes y, z \otimes w \rangle_C = \langle y, \langle x, z \rangle_B \cdot w \rangle_C$$

[Lan95, Proposition 4.5]. Modding out by the elements of length 0 and completing gives a right-Hilbert A – C bimodule $X \otimes_B Y$ in which $x \cdot b \otimes y = x \otimes b \cdot y$. We emphasize that it is not obvious that the action of $a \in A$ on $X \odot Y$ extends to $X \otimes_B Y$; this requires a complete positivity argument as in [Lan95, page 42]. Note that $X \otimes_B Y$ is full, since X and Y are:

$$\overline{\langle X \otimes_B Y, X \otimes_B Y \rangle_C} = \overline{\langle Y, \langle X, X \rangle_B \cdot Y \rangle_C} = \overline{\langle Y, B \cdot Y \rangle_C} = \overline{\langle Y, Y \rangle_C} = C.$$

We call $X \otimes_B Y$ the *balanced* or *internal tensor product*.

Proposition 2.4. *There is a category \mathcal{C} in which the objects are C^* -algebras, and in which the morphisms from A to B are the isomorphism classes of (full) right-Hilbert A – B bimodules. The composition of $[X]: A \rightarrow B$ with $[Y]: B \rightarrow C$ is the isomorphism class of the internal tensor product bimodule ${}_A(X \otimes_B Y)_C$; the identity morphism on A is the isomorphism class of the right-Hilbert bimodule ${}_A A_A$.*

Remark 2.5. In any category, $\text{Hom}(A, B)$ has to be a set for each pair of objects A and B . This is not obviously true in \mathcal{C} unless we limit the size of the bimodules involved. We can do this by considering only C^* -algebras and Hilbert modules with dense subsets whose cardinalities do not exceed a fixed large cardinal. For example, we could consider only separable C^* -algebras and bimodules. In practice, these issues should never present an unassailable problem, and we shall ignore them.

Proof of Proposition 2.4. We first claim that the composition of morphisms is well-defined. Suppose we have right-Hilbert bimodule isomorphisms $\Phi: {}_A X_B \rightarrow {}_A Z_B$ and $\Psi: {}_B Y_C \rightarrow {}_B W_C$. Then $\Phi \otimes \Psi: X \odot Y \rightarrow Z \otimes_B W$ is easily seen to preserve the actions and inner product, so extends to an isometric bimodule map $\Phi \otimes_B \Psi$. The map $\Phi^{-1} \otimes_B \Psi^{-1}$ is an inverse for $\Phi \otimes_B \Psi$, which is therefore a right-Hilbert A – C bimodule isomorphism of $X \otimes_B Y$ onto $Z \otimes_B W$.

Next we establish that composition of morphisms in \mathcal{C} is associative; it suffices to show that $X \otimes_B (Y \otimes_C Z)$ and $(X \otimes_B Y) \otimes_C Z$ are isomorphic for any right-Hilbert bimodules ${}_A X_B$, ${}_B Y_C$, and ${}_C Z_D$. But straightforward calculations show that the usual linear isomorphism of $X \odot (Y \odot Z)$ onto $(X \odot Y) \odot Z$ respects the module actions and right inner products, so extends to the desired isomorphism.

Finally, note that the maps $a \otimes x \mapsto a \cdot x$ and $y \otimes a \mapsto y \cdot a$ extend to isomorphisms $A \otimes_A X \cong X$ and $Y \otimes_A A \cong Y$ for any right-Hilbert bimodules ${}_A X_B$ and ${}_B Y_A$. Hence $[_A A_A]$ is an identity morphism from A to A in \mathcal{C} . \square

It is implicit in Connes' definition of Morita equivalence ([Con94, p. 155]) that a morphism $[X]$ is invertible in \mathcal{C} precisely when X is an imprimitivity bimodule. This seemed obvious to us at first but we found it surprisingly hard to prove.

By saying that a right-Hilbert bimodule ${}_AX_B$ is an imprimitivity bimodule we mean that X is also a (full) left-Hilbert A -module in such a way that

$${}_A\langle x, y \rangle \cdot z = x \cdot \langle y, z \rangle_B \quad \text{and} \quad {}_A\langle x \cdot b, y \rangle = {}_A\langle x, y \cdot b^* \rangle.$$

Equivalently, ${}_AX_B$ is an imprimitivity bimodule if and only if the canonical map $\kappa: A \rightarrow \mathcal{K}(X_B)$ induced by the left action of A on X_B is an isomorphism of A onto the algebra $\mathcal{K}(X_B)$ of compact operators on X_B [RW98, Proposition 3.8].

Proposition 2.6. *Suppose ${}_AX_B$ is a right-Hilbert bimodule. Then $[X]$ is an isomorphism in the category \mathcal{C} if and only if X is an A - B imprimitivity bimodule.*

One direction of Proposition 2.6 is easy: if ${}_AX_B$ is an imprimitivity bimodule, and ${}_B\tilde{X}_A$ is its reverse bimodule, then the maps $x \otimes \tilde{y} \mapsto {}_A\langle x, y \rangle$ and $\tilde{x} \otimes y \mapsto \langle x, y \rangle_B$ are isomorphisms of $X \otimes_B \tilde{X}$ onto ${}_AA$ and $\tilde{X} \otimes_A X$ onto ${}_BB$, respectively (see [RW98, Proposition 3.28]), and hence $[\tilde{X}]$ is an inverse for $[X]$ in \mathcal{C} .

For the other direction we want to use a representation-theoretic argument, so we briefly discuss how right-Hilbert bimodules can be represented by bounded operators. These ideas are fairly standard: the analogues for imprimitivity bimodules appear in [ER95, Section 2].

Definition 2.7. A *representation* (π_A, π_X, π_B) of a right-Hilbert bimodule ${}_AX_B$ is a triple consisting of nondegenerate representations π_A and π_B of A and B on Hilbert spaces \mathcal{H}_A and \mathcal{H}_B , respectively, together with a linear map $\pi_X: X \rightarrow \mathcal{B}(\mathcal{H}_B, \mathcal{H}_A)$ such that

- (i) $\pi_X(a \cdot x) = \pi_A(a)\pi_X(x)$,
- (ii) $\pi_X(x \cdot b) = \pi_X(x)\pi_B(b)$, and
- (iii) $\pi_B(\langle x, y \rangle_B) = \pi_X(x)^*\pi_X(y)$

for all $a \in A$, $x, y \in X$, and $b \in B$. (Again, property (ii) follows from property (iii), but we include it so the definition explicitly says that π preserves the Hilbert-module structure of X .)

Example 2.8. Suppose ${}_AX_B$ is a right-Hilbert bimodule and π_B is any nondegenerate representation of B on a Hilbert space \mathcal{H}_B . The *induced representation* $\pi_A = X\text{-Ind } \pi_B$ of A acts in $\mathcal{H}_A = X \otimes_B \mathcal{H}_B$ according to $\pi_A(a)(x \otimes \xi) = (a \cdot x) \otimes \xi$ [RW98, Proposition 2.66]. We claim that there are bounded operators $\pi_X(x) \in \mathcal{L}(\mathcal{H}_B, \mathcal{H}_A)$ such that $\pi_X(x)\xi = x \otimes \xi$, and that (π_A, π_X, π_B) is then a nondegenerate representation of ${}_AX_B$ on $(\mathcal{H}_A, \mathcal{H}_B)$. Indeed, it is easy to check that $\pi_X(x)$ is bounded:

$$\|\pi_X(x)\xi\|^2 = (x \otimes \xi \mid x \otimes \xi) = (\pi_B(\langle x, x \rangle_B)\xi \mid \xi) \leq \|\pi_B(\langle x, x \rangle_B)\| \|\xi\|^2;$$

the relation $\pi_A(a)\pi_X(x) = \pi_X(a \cdot x)$ is immediate from the definition of π_A ; the relation $\pi_X(x)\pi_B(b) = \pi_X(x \cdot b)$ follows from the balancing property $(x \cdot b) \otimes \xi = x \otimes \pi_B(b)\xi$ of the tensor product; and the definition of the inner product on $X \otimes_B \mathcal{H}_B$ implies that $\pi_B(\langle x, y \rangle_B) = \pi_X(x)^*\pi_X(y)$.

In this example, the representation π_A is by definition the restriction of a representation $\pi_{\mathcal{L}}$ of the algebra $\mathcal{L}(X_B)$ of adjointable operators on X (strictly speaking, $\pi_A = \pi_{\mathcal{L}} \circ \kappa$). The following observation will be useful in the proof of Proposition 2.6.

Lemma 2.9. *Let X_B be a right Hilbert module, let π_B be a faithful representation of B on \mathcal{H}_B , and consider the representation $(\pi_{\mathcal{L}}, \pi_X, \pi_B)$ of the right-Hilbert bimodule ${}_{\mathcal{L}(X)}X_B$ discussed in Example 2.8. Then $\pi_{\mathcal{L}}$ is faithful, and*

$$\pi_{\mathcal{L}}(\mathcal{K}(X_B)) = \pi_X(X)\pi_X(X)^* := \overline{\text{span}}\{\pi_X(x)\pi_X(y)^* \mid x, y \in X\}.$$

Proof. By definition, $\mathcal{K}(X_B)$ is the closed span of the rank-one operators $\{\Theta_{x,y} \mid x, y \in X\}$ [RW98, Definition 2.24], so it is enough to show that $\pi_{\mathcal{L}}(\Theta_{x,y}) = \pi_X(x)\pi_X(y)^*$. We can verify this by applying both sides to vectors of the form $x \otimes \xi$, which densely span $X \otimes_B \mathcal{H}_B$. We then have

$$\begin{aligned} \pi_{\mathcal{L}}(\Theta_{x,y})(z \otimes \xi) &= \Theta_{x,y}(z) \otimes \xi = x \cdot \langle y, z \rangle_B \otimes \xi = x \otimes \pi(\langle y, z \rangle_B)\xi \\ &= x \otimes \pi_X(y)^*\pi_X(z)\xi = \pi_X(x)(\pi_X(y)^*\pi_X(z)\xi) \\ &= (\pi_X(x)\pi_X(y)^*)(z \otimes \xi), \end{aligned}$$

as required.

Since π_B is faithful and X is a $\mathcal{K}(X_B)$ – B imprimitivity bimodule, the induced representation $\pi_{\mathcal{K}}$ is faithful (this follows from the Rieffel correspondence [RW98, Proposition 3.24]), and hence so is its extension $\pi_{\mathcal{L}}$ to $\mathcal{L}(X_B) = M(\mathcal{K}(X_B))$. \square

Proof of Proposition 2.6. We have to show that if there is a right-Hilbert bimodule ${}_BY_A$ such that $X \otimes_B Y \cong {}_AA_A$ and $Y \otimes_A X \cong {}_BB_B$ as right-Hilbert bimodules, then X is an A – B imprimitivity bimodule (and in fact Y will then be isomorphic to \tilde{X}). We choose a faithful nondegenerate representation π_B of B on \mathcal{H}_B , and use the inducing construction of Example 2.8 to give a representation (π_A, π_X, π_B) of ${}_AX_B$ on $(\mathcal{H}_B, \mathcal{H}_A)$ with $\mathcal{H}_A = X \otimes_B \mathcal{H}_B$. We shall prove the result by showing that the presence of Y implies that π_A is an isomorphism of A onto the image $\overline{\text{span}}\{\pi_X(x)\pi_X(y)^*\}$ of the algebra $\mathcal{K}(X_B)$ of compact operators.

We begin by noting that because A acts faithfully on itself and ${}_AA_A \cong X \otimes_B Y$, A must act faithfully on X : if $a \cdot x = 0$ for all x , then $a \cdot (x \otimes y) = 0$ for all y , whence $ab = 0$ for all $b \in A$, and $a = 0$. Thus $\kappa: A \rightarrow \mathcal{L}(X_B)$ is injective, and the injectivity of $\pi_{\mathcal{L}}$ implies that $\pi_A = \pi_{\mathcal{L}} \circ \kappa$ is faithful. To show that π_A has the right image, we construct a representation (π_B, σ, π_A) of Y on $(\mathcal{H}_A, \mathcal{H}_B)$.

Let $\rho: Y \rightarrow \mathcal{B}(\mathcal{H}_A, Y \otimes_A \mathcal{H}_A)$ be the representation induced from π_A , given as usual by $\rho(y)\zeta = y \otimes \zeta$. By assumption, there exists a right-Hilbert bimodule isomorphism $\Psi: Y \otimes_A X \rightarrow B$, which we use to define a unitary $U: Y \otimes_A \mathcal{H}_A = Y \otimes_A X \otimes_B \mathcal{H}_B \rightarrow \mathcal{H}_B$ by

$$U(y \otimes x \otimes \xi) = \pi_B(\Psi(y \otimes x))\xi.$$

Now we define

$$\sigma(y) = U \circ \rho(y).$$

For $y \in Y$, $b \in B$, and $x \otimes \xi \in \mathcal{H}_A$ we get

$$\begin{aligned}\sigma(b \cdot y)(x \otimes \xi) &= U(b \cdot y \otimes x \otimes \xi) = \pi_B(\Psi(b \cdot y \otimes x))\xi \\ &= \pi_B(b)\pi_B(\Psi(y \otimes x))\xi = \pi_B(b)U(y \otimes x \otimes \xi) \\ &= \pi_B(b)\sigma(y)(x \otimes \xi),\end{aligned}$$

and we also have

$$\sigma(y)^*\sigma(z) = \rho(y)^*U^*U\rho(z) = \rho(y)^*\rho(z) = \pi_A(\langle y, z \rangle_A)$$

for all $y, z \in Y$, since ρ has right coefficient map π_A . Thus (π_B, σ, π_A) is indeed a right-Hilbert bimodule representation of Y . In particular, we have $\pi_A(A) = \sigma(Y)^*\sigma(Y)$.

It only remains to show that $\sigma(Y)^* = \pi_X(X)$. For this we observe that for $y \in Y$, $x \in X$, and $\xi \in \mathcal{H}_B$ we have

$$\sigma(y)\pi_X(x)\xi = \sigma(y)(x \otimes \xi) = \pi_B(\Psi(y \otimes x))\xi,$$

hence $\sigma(Y)\pi_X(X) = \pi_B(\Psi(Y \otimes_A X)) = \pi_B(B)$. Using $\pi_A(A) = \sigma(Y)^*\sigma(Y)$, it now follows that

$$\begin{aligned}\pi_X(X) &= \pi_X(A \cdot X) = \pi_A(A)\pi_X(X) = \sigma(Y)^*\sigma(Y)\pi_X(X) \\ &= \sigma(Y)^*\pi_B(B) = \sigma(Y \cdot B)^* = \sigma(Y)^*.\end{aligned}$$

This completes the proof of Proposition 2.6. \square

Proposition 2.10. *For every morphism $[{}_AX_B]$ in \mathcal{C} , there exists a C^* -algebra C , a nondegenerate homomorphism $\varphi: A \rightarrow M(C)$, and an imprimitivity bimodule ${}_CY_B$ such that*

$$[{}_AX_B] = [{}_CY_B] \circ [\varphi] \quad \text{in } \mathcal{C}.$$

In other words, such that ${}_AX_B \cong {}_A(C \otimes_C Y)_B$ as a right-Hilbert bimodule.

Proof. Let $C = \mathcal{K}(X_B)$ and let $Y = X$ viewed as a C - B imprimitivity bimodule (see [RW98, Proposition 3.8]). Since $\mathcal{L}(X_B) \cong M(\mathcal{K}(X_B))$ ([RW98, Corollary 2.54]), we can view the canonical nondegenerate homomorphism $\kappa: A \rightarrow \mathcal{L}(X_B)$ as a nondegenerate homomorphism $\varphi: A \rightarrow M(\mathcal{K}(X_B))$. The map $c \otimes y \mapsto c \cdot y$ extends to an isomorphism of $C \otimes_C Y$ onto X . \square

3. CROSSED-PRODUCT FUNCTORS

Our next goal is to formalize the idea that assignments like $(A, \alpha) \mapsto A \rtimes_\alpha G$ are functors into \mathcal{C} ; this requires that we construct an equivariant category $\mathcal{A}(G)$ in which the objects are dynamical systems (A, G, α) .

Definition 3.1. Let G be a locally compact group, let ${}_AX_B$ be a right-Hilbert bimodule, and let α and β be (strongly continuous) actions of G on A and B . An (α, β) -compatible action of G on X is a homomorphism γ of G into the group of invertible linear transformations on X such that

$$(i) \quad \gamma_s(a \cdot x) = \alpha_s(a) \cdot \gamma_s(x)$$

- (ii) $\gamma_s(x \cdot b) = \gamma_s(x) \cdot \beta_s(b)$
- (iii) $\langle \gamma_s(x), \gamma_s(y) \rangle_B = \beta_s(\langle x, y \rangle_B)$

for each $s \in G$, $a \in A$, $x, y \in X$, and $b \in B$; and such that each map $s \mapsto \gamma_s(x)$ is continuous from G into X . (As usual, (ii) follows from (iii).) Two (α, β) -compatible actions γ and η on X and Y are *equivariantly isomorphic* if there exists an isomorphism Φ of X onto Y such that $\Phi(\gamma_s(x)) = \eta_s(\Phi(x))$ for each $s \in G$ and $x \in X$.

Example 3.2. If (A, α) and (B, β) are actions of G , and $\varphi: A \rightarrow M(B)$ is equivariant in the sense that $\varphi(\alpha_s(a)) = \beta_s(\varphi(a))$, then β is an (α, β) compatible action of G on the bimodule ${}_A B_B$ of Example 2.2.

Proposition 3.3. *Let G be a locally compact group.*

- (i) *There is a category $\mathcal{A}(G)$ in which the objects are C^* -algebras with actions of G , and in which the morphisms from (A, α) to (B, β) are the equivariant isomorphism classes of (full) right-Hilbert A - B bimodules with (α, β) -compatible actions of G . The composition of $[X, \gamma]: (A, \alpha) \rightarrow (B, \beta)$ with $[Y, \eta]: (B, \beta) \rightarrow (C, \epsilon)$ is the isomorphism class of the tensor product action $(X \otimes_B Y, \gamma \otimes_B \eta)$; the identity morphism on (A, α) is $[(A, \alpha)_{(A, \alpha)}]$.*
- (ii) *$[(A, \alpha)(X, \gamma)_{(B, \beta)}]$ is an isomorphism in $\mathcal{A}(G)$ if and only if ${}_A X_B$ is an imprimitivity bimodule and $\alpha_s(\langle x, y \rangle) = {}_A \langle \gamma_s(x), \gamma_s(y) \rangle$.*
- (iii) *For every morphism $[(A, \alpha)(X, \gamma)_{(B, \beta)}]$ in $\mathcal{A}(G)$, there exists an isomorphism $[(C, \epsilon)(Y, \eta)_{(B, \beta)}]$ and an α - ϵ equivariant nondegenerate homomorphism $\varphi: A \rightarrow M(C)$ such that*

$$[X, \gamma] = [Y, \eta] \circ [\varphi, \epsilon] \quad \text{in } \mathcal{A}(G).$$

Proof. Adding actions to Propositions 2.4, 2.6, and 2.10 is routine, except possibly in statement (iii). In the proof of Proposition 2.10, we took $C = \mathcal{K}(X_B)$ and $\varphi = \kappa: A \rightarrow \mathcal{L}(X_B) = M(C)$, so we need to show that γ induces an action on $\mathcal{K}(X_B)$. But for each $T \in \mathcal{L}(X_B)$, $\epsilon_s(T): x \mapsto \gamma_s(T(\gamma_s^{-1}(x)))$ is an adjointable operator with $\epsilon_s(T)^* = \epsilon_s(T^*)$, and in fact $T \mapsto \epsilon_s(T)$ is an automorphism of $\mathcal{L}(X_B)$. A quick calculation shows that

$$(3.1) \quad \epsilon_s(\langle x, y \rangle) = \langle \gamma_s(x), \gamma_s(y) \rangle,$$

so ϵ_s restricts to an automorphism of $\mathcal{K}(X_B)$. Since G acts continuously on X , ϵ gives an action of G on \mathcal{K} , and (3.1) says that $[(C, \epsilon)(X, \gamma)_{(B, \beta)}]$ is an isomorphism in $\mathcal{A}(G)$. Now recall that $\kappa(\alpha_s(a))(x) = \alpha_s(a) \cdot x = \gamma_s(a \cdot \gamma_s^{-1}(x))$ to see that φ is α - ϵ equivariant; it is then easy to check that the isomorphism $c \otimes y \mapsto c \cdot y$ of $C \otimes_C Y$ onto X is $\epsilon \otimes \eta$ - γ equivariant. \square

To construct the crossed product of a system (A, G, α) , we begin with the vector space $C_c(G, A)$ of continuous functions $f: G \rightarrow A$ of compact support. This is a $*$ -algebra with the operations

$$f * g(s) = \int_G f(t) \alpha_t(g(t^{-1}s)) dt \quad \text{and} \quad f^*(s) = \alpha_s(f(s^{-1})^*) \Delta_G(s)^{-1}.$$

A covariant representation of (A, G, α) on \mathcal{H} is a pair (π, U) consisting of a nondegenerate representation $\pi: A \rightarrow \mathcal{B}(\mathcal{H})$ and a unitary representation $U: G \rightarrow U(\mathcal{H})$ such that $\pi(\alpha_s(a)) = U_s \pi(a) U_s^*$ for $s \in G$ and $a \in A$. Routine calculations show that there is a $*$ -representation $\pi \times U$ of $C_c(G, A)$ on \mathcal{H} , called the integrated form of (π, U) , such that

$$\pi \times U(f)\xi = \int_G \pi(f(s))U_s \xi \, ds$$

for all $\xi \in \mathcal{H}$. (Inserting the vector ξ is technically helpful because it makes the integrand norm-continuous.) Since $\|\pi \times U(f)\| \leq \int_G \|f(s)\| \, ds$ for every f , we can define a semi-norm on $C_c(G, A)$ by ²

$$\|f\|_* = \sup\{\|\pi \times U(f)\| : (\pi, U) \text{ is a covariant representation of } (A, G, \alpha)\};$$

it is a norm because the regular representation $\tilde{\pi} \times \lambda$ on $L^2(G, \mathcal{H}_\pi)$ is faithful on $C_c(G, A)$ if π is faithful on A .

We now define the crossed product $A \rtimes_\alpha G$ to be the completion of $C_c(G, A)$ in the norm $\|\cdot\|_*$. (For this to be a construction of a particular C^* -algebra rather than an isomorphism class of C^* -algebras, we must be clear that forming completions is a construction, but this can be achieved by defining the completion of a normed space X to be the closure of X in its double dual.) Notice that every representation $\pi \times U$ of $C_c(G, A)$ is by definition continuous for $\|\cdot\|_*$, and hence extends to a representation of $A \rtimes_\alpha G$, which we continue to denote by $\pi \times U$.

Remark 3.4. In [Rae88], a crossed product for the system (A, G, α) is any C^* -algebra B equipped with a nondegenerate homomorphism $i_A: A \rightarrow M(B)$ and a strictly continuous homomorphism $i_G: G \rightarrow UM(B)$ satisfying

- (a) (i_A, i_G) is covariant, in the sense that $i_A(\alpha_s(a)) = i_G(s)i_A(a)i_G(s)^*$ for all $a \in A$ and $s \in G$;
- (b) if (π, U) is a covariant representation, there is a nondegenerate representation $\varphi = \varphi_{\pi, U}$ of B such that $\pi = \varphi \circ i_A$ and $U = \varphi \circ i_G$; and
- (c) the elements $i_A \times i_G(f) = \int_G i_A(f(s))i_G(s) \, ds$ for $f \in C_c(G, A)$ span a dense subspace of B .

From our present perspective, we can view these axioms as properties which allow us to identify isomorphic copies of $A \rtimes_\alpha G$. To be more precise, if (B, i_A, i_G) is a crossed product in the sense of [Rae88], then we claim that $i_A \times i_G$ extends to an isomorphism of $A \rtimes_\alpha G$ onto B .

To see this, first represent B on Hilbert space, so that by (a), (i_A, i_G) becomes a covariant representation of (A, G, α) . It follows from the definition of $\|\cdot\|_*$ that $i_A \times i_G$ extends to a representation of $A \rtimes_\alpha G$, and then (c) implies that $i_A \times i_G$ maps

² There is ostensibly a set-theoretic problem here since it is not obvious that we are taking the supremum over a well-defined set of real numbers. But the numbers of the form $\|\pi \times U(f)\|$ do form a subclass of \mathbf{R} , and every subclass of a set is again a set ([Men64, Proposition 4.6]).

$A \times_\alpha G$ onto (the represented copy of) B . Now choose any covariant representation (π, U) of (A, G, α) . Then by (b), $\pi \times U(f) = \varphi_{\pi, U}(i_A \times i_G(f))$ for $f \in C_c(G, A)$, and hence for all $f \in A \times_\alpha G$. But this implies $\|\pi \times U(f)\| \leq \|i_A \times i_G(f)\|$, and hence that $\|f\|_* \leq \|i_A \times i_G(f)\|$. Thus $i_A \times i_G$ is injective, which proves the claim.

We next seek to define crossed products of right-Hilbert bimodules. There are several possible approaches; for example, we could deduce much of the following proposition from the construction of [Com84, Section 5] for imprimitivity bimodules by first factoring the right-Hilbert bimodule as in Proposition 3.3(iii). We have opted for a direct treatment partly because it is more elementary, and partly because we feel that the details should be available. This is also the approach used by Kasparov to construct crossed-product Hilbert modules for the study of the KK -theory of group C^* -algebras in [Kas95][§6, Definition 1].

Proposition 3.5. *Let γ be an (α, β) -compatible action of G on a right-Hilbert bimodule ${}_A X_B$. There exists a right-Hilbert $(A \times_\alpha G) - (B \times_\beta G)$ bimodule $X \times_\gamma G$, which contains $C_c(G, X)$ as a dense subspace, and which satisfies*

$$(3.2) \quad f \cdot h(s) = \int_G f(t) \cdot \gamma_t(h(t^{-1}s)) dt,$$

$$(3.3) \quad h \cdot g(s) = \int_G h(t) \cdot \beta_t(g(t^{-1}s)) dt, \quad \text{and}$$

$$(3.4) \quad \langle h, k \rangle_{B \times_\beta G}(s) = \int_G \beta_{t^{-1}}(\langle h(t), k(ts) \rangle_B) dt$$

for $f \in C_c(G, A)$, $h, k \in C_c(G, X)$, and $g \in C_c(G, B)$.

Lemma 3.6. *Let $\pi \times U$ be a representation of $B \times_\beta G$ on a Hilbert space \mathcal{H} . Then for each $h, k \in C_c(G, X)$ and $\xi, \eta \in \mathcal{H}$, (3.4) satisfies*

$$(\pi \times U(\langle k, h \rangle_{B \times_\beta G})\xi \mid \eta) = (\omega_{h, \xi} \mid \omega_{k, \eta}),$$

where $\omega_{h, \xi} = \int_G h(s) \otimes U_s \xi ds \in X \otimes_B \mathcal{H}$.

Proof. We have

$$\begin{aligned}
(\pi \times U(\langle k, h \rangle_{B \times G}) \xi \mid \eta) &= \left(\int_G \pi(\langle k, h \rangle_{B \times G}(s)) U_s \xi \, ds \mid \eta \right) \\
&= \left(\int_G \pi \left(\int_G \beta_{t^{-1}}(\langle k(t), h(ts) \rangle_B) \, dt \right) U_s \xi \, ds \mid \eta \right) \\
&= \int_G \int_G (U_t^* \pi(\langle k(t), h(ts) \rangle_B) U_{ts} \xi \mid \eta) \, ds \, dt \\
&= \int_G \int_G (\pi(\langle k(t), h(s) \rangle_B) U_s \xi \mid U_t \eta) \, ds \, dt \\
&= \int_G \int_G (h(s) \otimes U_s \xi \mid k(t) \otimes U_t \eta) \, ds \, dt \\
&= \left(\int_G h(s) \otimes U_s \xi \, ds \mid \int_G k(t) \otimes U_t \eta \, dt \right) \\
&= (\omega_{h, \xi} \mid \omega_{k, \eta}).
\end{aligned}$$

□

Proof of Proposition 3.5. We begin by showing that $C_c(G, X)$ can be completed to give a Hilbert $B \times_\beta G$ -module $X \times_\gamma G$ satisfying (3.3) and (3.4). Straightforward calculations show that (3.3) and (3.4) make $C_c(G, X)$ into a pre-inner product $B \times_\beta G$ -module, so we need only verify that the sesquilinear form of (3.4) is positive definite.

To do so, fix, for the remainder of the proof, a faithful representation $\pi \times U$ of $B \times_\beta G$ on a Hilbert space \mathcal{H} . Then for each $\xi \in \mathcal{H}$, Lemma 3.6 gives

$$(\pi \times U(\langle h, h \rangle_{B \times G}) \xi \mid \xi) = (\omega \mid \omega) \geq 0,$$

which shows positivity. For definiteness, suppose $h \in C_c(G, X)$ satisfies $\langle h, h \rangle_{B \times G} = 0$. Then Lemma 3.6 gives $(\pi \times U(\langle h, h \rangle_{B \times G}) \xi \mid \xi) = (\omega \mid \omega) = 0$ for each $\xi \in \mathcal{H}$, so that $\omega = \int_G h(s) \otimes U_s \xi \, ds = 0$ for each ξ . Thus,

$$\left(\int_G h(s) \otimes U_s \xi \, ds \mid x \otimes \zeta \right) = \left(\int_G \pi(\langle x, h(s) \rangle_B) U_s \xi \, ds \mid \zeta \right) = (\pi \times U(g_x) \xi \mid \zeta) = 0$$

for each $\xi, \zeta \in \mathcal{H}$ and $x \in X$, where $g_x(s) = \langle x, h(s) \rangle_B$ defines $g_x \in C_c(G, X)$. It follows that $g_x = 0$ for all $x \in X$, whence $h = 0$ in $C_c(G, X)$.

We next show that the $(B \times_\beta G)$ -valued inner product on $X \times_\gamma G$ is full. Since $C_c(G, B)$ has an approximate identity for $B \times_\beta G$, since $\langle X, X \rangle_B$ is dense in B , and since B acts nondegenerately on $B \times_\beta G$, functions of the form $f^* \langle x, y \rangle_B g$, where $f, g \in C_c(G, B)$ and $x, y \in X$, span a dense subspace of $B \times_\beta G$. Now letting $(x \cdot f)(s) = x \cdot (f(s))$ define $x \cdot f \in C_c(G, X)$, a straightforward calculation shows that $\langle x \cdot f, y \cdot g \rangle_{B \times G} = f^* \langle x, y \rangle_B g$, so that $\langle C_c(G, X), C_c(G, X) \rangle_{B \times G}$ is dense in $B \times_\beta G$.

We now claim that (3.2) makes $X \times_\gamma G$ into a right-Hilbert $(A \times_\alpha G)$ – $(B \times_\beta G)$ bimodule. Again, checking the algebraic conditions of Definition 2.1 at the level of

C_c -functions is routine; we need to show that $\|f \cdot h\| \leq \|f\| \|h\|$ for $f \in C_c(G, A)$ and $h \in C_c(G, X)$ to see that this extends to an action of $A \times_\alpha G$ on $X \times_\gamma G$.

To show this, we begin by defining actions of A and G on $C_c(G, X)$ by

$$(a \cdot h)(s) = a \cdot (h(s)) \quad \text{and} \quad (t \cdot h)(s) = \gamma_t(h(t^{-1}s)).$$

Now we use these actions to define

$$\rho(a)(h \otimes \xi) = a \cdot h \otimes \xi \quad \text{and} \quad V_t(h \otimes \xi) = t \cdot h \otimes \xi;$$

we claim that $\rho(a)$ is bounded on $C_c(G, X) \odot \mathcal{H}$, and hence extends to an operator on $(X \times_\gamma G) \otimes_{B \times_\beta G} \mathcal{H}$. Again using Lemma 3.6, and writing ω_i for ω_{h_i, ξ_i} , we have

$$\begin{aligned} \left\| \rho(a) \left(\sum_{i=1}^n h_i \otimes \xi_i \right) \right\|^2 &= \left(a \cdot \sum_{i=1}^n h_i \otimes \xi_i \mid a \cdot \sum_{j=1}^n h_j \otimes \xi_j \right) \\ &= \sum_{i,j=1}^n (a \cdot h_i \otimes \xi_i \mid a \cdot h_j \otimes \xi_j) \\ &= \sum_{i,j=1}^n (\pi \times U(\langle a \cdot h_j, a \cdot h_i \rangle_{B \times G}) \xi_i \mid \xi_j) \\ &= \sum_{i,j=1}^n \left(\int_G a \cdot h_i(s) \otimes U_s \xi_i \, ds \mid \int_G a \cdot h_j(s) \otimes U_s \xi_j \, ds \right) \\ &= \sum_{i,j=1}^n (a \cdot \omega_i \mid a \cdot \omega_j) \\ &= \left(a \cdot \sum_{i=1}^n \omega_i \mid a \cdot \sum_{j=1}^n \omega_j \right) \\ &\leq \|a\|^2 \sum_{i,j=1}^n (\omega_i \mid \omega_j) \\ &= \|a\|^2 \sum_{i,j=1}^n (\pi \times U(\langle h_j, h_i \rangle_{B \times G}) \xi_i \mid \xi_j) \\ &= \|a\|^2 \left(\sum_{i=1}^n h_i \otimes \xi_i \mid \sum_{j=1}^n h_j \otimes \xi_j \right) \\ &= \|a\|^2 \left\| \sum_{i=1}^n h_i \otimes \xi_i \right\|^2, \end{aligned}$$

where the inequality holds because A acts boundedly on $X \otimes_B \mathcal{H}$ via the induced representation X -Ind π .

It is straightforward to check that each V_t is unitary, and then that (ρ, V) is co-variant for (A, G, α) ; ρ is nondegenerate because A acts nondegenerately on X , so we get a nondegenerate representation $\rho \times V$ of $A \times_\alpha G$ on $(X \times_\gamma G) \otimes_{B \times G} \mathcal{H}$.

Now for $f \in C_c(G, A)$, $h \in C_c(G, X)$, and $\xi \in \mathcal{H}$, we have

$$\rho \times V(f)(h \otimes \xi) = \int_G \rho(f(t))V_t(h \otimes \xi) dt = \left(\int_G f(t)t \cdot h dt \right) \otimes \xi = f \cdot h \otimes \xi,$$

so

$$\begin{aligned} (\pi \times U(\langle f \cdot h, f \cdot h \rangle_{B \times G})\xi \mid \xi) &= (f \cdot h \otimes \xi \mid f \cdot h \otimes \xi) \\ &= (\rho \times V(f)(h \otimes \xi) \mid \rho \times V(f)(h \otimes \xi)) \\ &\leq \|f\|^2 (h \otimes \xi \mid h \otimes \xi) \\ &= (\pi \times U(\|f\|^2 \langle h, h \rangle_{B \times G})\xi \mid \xi). \end{aligned}$$

It follows that $\|f \cdot h\| \leq \|f\| \|h\|$.

Finally, to see that the action of $A \times_\alpha G$ on $X \times_\gamma G$ is nondegenerate, note that for $h \in C_c(G, X)$, the Cauchy-Schwarz inequality gives

$$\begin{aligned} \|h\|^2 &= \|\langle h, h \rangle_{B \times G}\| \\ &\leq \int_G \|\langle h, h \rangle_{B \times G}(s)\| ds \\ &\leq \int_G \int_G \|\beta_{t^{-1}}(\langle h(t), h(ts) \rangle_B)\| dt ds \\ &\leq \int_G \int_G \|h(t)\| \|h(ts)\| dt ds \\ &\leq \left(\int_G \|h(t)\| dt \right)^2, \end{aligned}$$

so $\|h\| \leq \|h\|_1$. Now standard arguments show that we can choose $f \in C_c(G, A)$ to make $\|h - f \cdot h\|_1$ arbitrarily small: take f of the form $s \mapsto a\chi(s)$ as a runs through an approximate identity for A and χ runs through an approximate identity for $C_c(G)$. \square

Example 3.7. Suppose (A, α) and (B, β) are objects in $\mathcal{A}(G)$, and $\varphi : A \rightarrow M(B)$ is a nondegenerate homomorphism such that $\varphi \circ \alpha_s = \beta_s \circ \varphi$. Then the crossed-product morphism ${}_A B_B \times G$ is the completion of $C_c(G, B)$ in the norm coming from $B \times_\beta G$, and hence is precisely $(B \times_\beta G)_{B \times_\beta G}$; the left action is given by a nondegenerate homomorphism $\varphi \times G : A \times_\alpha G \rightarrow M(B \times_\beta G)$. Equation 3.2 shows that for functions $f \in C_c(G, A)$ and $g \in C_c(G, B)$, we have

$$((\varphi \times G)(f)g)(s) = \int_G \varphi(f(t))\beta_t(g(t^{-1}s)) dt.$$

Proposition 3.8. *The maps defined by*

$$(A, \alpha) \mapsto A \times_\alpha G \quad \text{and} \quad [(A, \alpha)(X, \gamma)_{(B, \beta)}] \mapsto [A \times_\alpha G(X \times_\gamma G)_{B \times G}]$$

give a functor from $\mathcal{A}(G)$ to \mathcal{C} .

Proof. We first show that the map on morphisms is well-defined. Suppose $\varphi: X \rightarrow Y$ is an isomorphism of right-Hilbert A – B bimodules which is equivariant for (α, β) -compatible actions γ and η of G . Then $\Phi(h)(s) = \varphi(h(s))$ is easily seen to give a bijective map $\Phi: C_c(G, X) \rightarrow C_c(G, Y)$ which respects the right-Hilbert bimodule structures (Equations 3.2–3.4) and hence extends to a right-Hilbert $(A \times_\alpha G)$ – $(B \times_\beta G)$ bimodule isomorphism of $X \times_\gamma G$ onto $Y \times_\eta G$. Example 3.7 shows that identity morphisms go to identity morphisms.

It only remains to see that the assignment $[X, \gamma] \mapsto [X \times_\gamma G]$ respects composition of morphisms; that is, if $({}_A X_B, \gamma)$ is (α, β) -compatible and $({}_B Y_C, \eta)$ is (β, ϵ) -compatible, we need to show that

$$(X \times_\gamma G) \otimes_{B \times G} (Y \times_\eta G) \cong (X \otimes_B Y) \times_{\gamma \otimes \eta} G$$

as right-Hilbert $A \times_\alpha G$ – $C \times_\epsilon G$ bimodules.³

The rule

$$\Psi(h \otimes k)(s) = \int_G h(t) \otimes \eta_t(k(t^{-1}s)) dt$$

defines a linear map from $C_c(G, X) \odot C_c(G, Y)$ to $C_c(G, X \otimes_B Y)$ which preserves the pre-right-Hilbert bimodule structures. In order to see that Ψ extends to an isomorphism of the completions, we need only verify that Ψ has dense range for the inductive limit topology. For this, let $x \in X$ and $f \in C_c(G, B)$, and define $h \in C_c(G, X)$ by $h(s) = x \cdot f(s)$. Then for $k \in C_c(G, Y)$ we have

$$\begin{aligned} \Phi(h \otimes k)(s) &= x \otimes \int_G f(t) \cdot \eta_t(k(t^{-1}s)) dt \\ &= x \otimes (f \cdot k)(s). \end{aligned}$$

Now, we can approximate k by $f \cdot k$ in the inductive limit topology, and taking k of the form $k(s) = yg(s)$ for $y \in Y$ and $g \in C_c(G)$ we can thus approximate the function $s \mapsto (x \otimes y)g(s)$. But such functions have inductive-limit-dense span in $C_c(G, X \otimes_B Y)$.

□

4. NATURALITY IN GREEN'S IMPRIMITIVITY THEOREM

Suppose that α is an action of a locally compact group G on a C^* -algebra A , and H is a closed subgroup of G . Takesaki showed in [Tak67] how Mackey's construction of induced representations of groups could be modified to induce covariant representations of (A, H, α) to representations of (A, G, α) . Subsequently Green showed how the integrated forms of Takesaki's induced representations could be obtained using Rieffel's abstract induction by Hilbert bimodules [Gre78]. To construct his

³An analog of this result appears as Lemma 3.10 in [Kas88], in the context of not-necessarily-full Hilbert modules.

$(A \times_\alpha H)$ – $(A \times_\alpha G)$ bimodule $X_H^G(A, \alpha)$, Green completed the $C_c(G, A)$ – $C_c(H, A)$ bimodule $C_c(G, A)$ in which

$$\begin{aligned} f \cdot x(t) &= \int_G f(s) \alpha_s(x(s^{-1}t)) \Delta(s)^{1/2} ds \\ x \cdot g(t) &= \int_H x(th) \alpha_{th}(g(h^{-1})) \Delta(h)^{-1/2} dh \\ \langle x, y \rangle_{A \times_\alpha H}(h) &= \int_G \alpha_s(x(s^{-1})^* y(s^{-1}h)) \Delta(h)^{-1/2} ds \end{aligned}$$

for $x \in C_c(G, A)$, $f \in C_c(G, A) \subset A \times_\alpha G$, $g \in C_c(H, A)$. The representation of $A \times_\alpha G$ induced from a representation π of $A \times_\alpha H$ is then by definition the representation X_H^G –Ind π acting in $X_H^G \otimes_{A \times H} \mathcal{H}_\pi$.

Imprimitivity theorems tell us how to recognize induced representations. In Rieffel’s modern formulation, we recognize a representation which has been induced from H by the presence of an auxiliary representation of $C_0(G/H)$. In Green’s realization, this representation comes from the action of $C_0(G/H)$ on $C_c(G, A)$ given by

$$c \cdot x(t) = c(th)x(t),$$

and the crucial properties of this action are that it commutes with the left action of A on X_H^G and is covariant for the left action of G : more formally,

$$c \cdot (i_A(a) \cdot x) = i_A(a) \cdot (c \cdot x) \quad \text{and} \quad i_G(s) \cdot (c \cdot (i_G(s)^* \cdot x)) = \tau_s(c) \cdot x,$$

where we have extended the left action of $A \times_\alpha G$ on X_H^G to $M(A \times_\alpha G)$ and τ is the action of G on $C_0(G/H)$ given by $\tau_s(c)(t) = c(s^{-1}t)$. In other words, the left actions combine to give an action of $(A \otimes C_0(G/H)) \times_{\alpha \otimes \tau} G$ on $X_H^G(A, \alpha)$, and it is the content of Green’s Imprimitivity Theorem that X_H^G is an $((A \otimes C_0(G/H)) \times_{\alpha \otimes \tau} G)$ – $(A \times_\alpha H)$ imprimitivity bimodule. Note that the left action of $A \times_\alpha G$ on X_H^G factors through the left action of $(A \otimes C_0(G/H)) \times_{\alpha \otimes \tau} G$ via the canonical map of $A \times G$ into $M((A \otimes C_0(G/H)) \times_{\alpha \otimes \tau} G)$.

Theorem 4.1. *Suppose H is a closed subgroup of a locally compact group G . Then the assignment $(A, \alpha) \mapsto [X_H^G(A, \alpha)]$ is a natural transformation between the functors $(A, \alpha) \mapsto A \times_\alpha G$ and $(A, \alpha) \mapsto A \times_\alpha H$ from $\mathcal{A}(G)$ to \mathcal{C} , and a natural equivalence between the functors $(A, \alpha) \mapsto (A \otimes C_0(G/H)) \times_{\alpha \otimes \tau} G$ and $(A, \alpha) \mapsto A \times_\alpha H$ from $\mathcal{A}(G)$ to \mathcal{C} .*

Before we begin the proof, we point out that we have implicitly asserted in the statement of the theorem that the crossed products $A \times_\alpha H$ and $(A \otimes C_0(G/H)) \times_{\alpha \otimes \tau} G$ define functors on $\mathcal{A}(G)$. The first case is quite easy: we just need to check that restricting every action in sight from G to H is a functor from $\mathcal{A}(G)$ to $\mathcal{A}(H)$, and then compose this with the crossed-product functor from $\mathcal{A}(H)$ to \mathcal{C} . For the second, we need to prove that $(A, \alpha) \mapsto (A \otimes C_0(G/H), \alpha \otimes \tau)$ is a functor from $\mathcal{A}(G)$ to itself, or, more generally, that for any given action ϵ of G on a C^* -algebra C ,

$(A, \alpha) \mapsto (A \otimes_{\min} C, \alpha \otimes \epsilon)$ is a functor. This is non-trivial: a morphism $[{}_A X_B, \gamma]$ goes to a morphism $[X \otimes C, \gamma \otimes \epsilon]$ based on the external tensor product ${}_{A \otimes C}(X \otimes C)_{B \otimes C}$ of bimodules (see [RW98, Corollary 3.38]). But all the details are routine, so we shall omit them.

A natural transformation T between two functors $F, G : \mathcal{A} \rightarrow \mathcal{B}$ assigns to each object A of \mathcal{A} a morphism $T(A) : F(A) \rightarrow G(A)$ such that, for every morphism $\varphi : A \rightarrow B$ in \mathcal{A} , the diagram

$$\begin{array}{ccc} F(A) & \xrightarrow{T(A)} & G(A) \\ F(\varphi) \downarrow & & \downarrow G(\varphi) \\ F(B) & \xrightarrow{T(B)} & G(B) \end{array}$$

commutes in \mathcal{B} . The transformation T is a natural equivalence if $T(A)$ is an isomorphism for all objects A . Green's Theorem tells us that $[X_H^G(A, \alpha)]$ is a morphism from $A \times_{\alpha} G$ to $A \times_{\alpha} H$ and an isomorphism from $(A \otimes C_0(G/H)) \times_{\alpha \otimes \tau} G$ to $A \times_{\alpha} H$, so we need to show that for every morphism $[(A, \alpha)(X, \gamma)_{(B, \beta)}]$ in the category $\mathcal{A}(G)$, the diagrams

$$(4.1) \quad \begin{array}{ccc} A \times_{\alpha} G & \xrightarrow{X_H^G(A, \alpha)} & A \times_{\alpha} H \\ X \times_{\gamma} G \downarrow & & \downarrow X \times_{\gamma} H \\ B \times_{\beta} G & \xrightarrow{X_H^G(B, \beta)} & B \times_{\beta} H \end{array}$$

and

$$(4.2) \quad \begin{array}{ccc} (A \otimes C_0(G/H)) \times_{\alpha \otimes \tau} G & \xrightarrow{X_H^G(A, \alpha)} & A \times_{\alpha} H \\ (X \otimes C_0(G/H)) \times_{\gamma \otimes \tau} G \downarrow & & \downarrow X \times_{\gamma} H \\ (B \otimes C_0(G/H)) \times_{\beta \otimes \tau} G & \xrightarrow{X_H^G(B, \beta)} & B \times_{\beta} H \end{array}$$

commute in \mathcal{C} .

Proof of Theorem 4.1. We establish the commutativity of both (4.1) and (4.2) by factoring $[X, \gamma] = [Y, \eta] \circ [\varphi, \epsilon]$, as in Proposition 3.3(iii), where $[(C, \epsilon)(Y, \eta)_{(B, \beta)}]$ is an isomorphism in $\mathcal{A}(G)$ and $\varphi : A \rightarrow M(C)$ is an α - ϵ equivariant nondegenerate homomorphism. We shall prove the commutativity of (4.1) by showing that we have

two commutative diagrams

$$(4.3) \quad \begin{array}{ccc} A \times_{\alpha} G & \xrightarrow{X_H^G(A, \alpha)} & A \times_{\alpha} H \\ \varphi \times G \downarrow & & \downarrow \varphi \times H \\ C \times_{\epsilon} G & \xrightarrow{X_H^G(C, \epsilon)} & C \times_{\epsilon} H \end{array}$$

and

$$(4.4) \quad \begin{array}{ccc} C \times_{\epsilon} G & \xrightarrow{X_H^G(C, \epsilon)} & C \times_{\epsilon} H \\ Y \times_{\eta} G \downarrow & & \downarrow Y \times_{\eta} H \\ B \times_{\beta} G & \xrightarrow{X_H^G(B, \beta)} & B \times_{\beta} H. \end{array}$$

Because taking crossed products is a functor, this will then give

$$\begin{aligned} [X_H^G(B)] \circ [A \times G(X \times_{\gamma} G)_{B \times G}] &= [X_H^G(B)] \circ [C \times G(Y \times_{\eta} G)_{B \times G}] \circ [\varphi \times G] \\ &= [C \times H(Y \times_{\eta} H)_{B \times H}] \circ [\varphi \times H] \circ [X_H^G(A)] \\ &= [A \times H(X \times_{\gamma} H)_{B \times H}] \circ [X_H^G(A)], \end{aligned}$$

as required.

The commutativity of (4.3) amounts to:

Lemma 4.2. *Suppose (A, α) , (C, ϵ) are objects in $\mathcal{A}(G)$ and $\varphi : A \rightarrow M(C)$ is a nondegenerate homomorphism such that $\varphi \circ \alpha = \epsilon \circ \varphi$. Then*

$$X_H^G(A, \alpha) \otimes_{A \times H} (C \times_{\epsilon} H) \cong X_H^G(C, \epsilon) \cong (C \times_{\epsilon} G) \otimes_{C \times G} X_H^G(C, \epsilon)$$

as right-Hilbert $(A \times_{\alpha} G)-(C \times_{\epsilon} H)$ bimodules.

Proof. The second isomorphism is standard. For the first, define

$$\Psi : C_c(G, A) \odot C_c(H, C) \rightarrow C_c(G, C)$$

by

$$\Psi(x \otimes g)(t) = \int_H \varphi(x(th)) \epsilon_{th}(g(h^{-1})) \Delta(h)^{-1/2} dh.$$

The usual change-of-variables arguments show that Ψ is right $C_c(H, C)$ -linear (though strictly speaking it is not necessary to prove this). The map $\varphi \times H$ is given by

$$(4.5) \quad (\varphi \times H(f)g)(h) = \int_H \varphi(f(k)) \epsilon_k(g(k^{-1}h)) dk;$$

using this identity, some convoluted calculations involving several changes of variables show that

$$\langle \Psi(x \otimes g), \Psi(y \otimes f) \rangle_{C_c(H, C)} = (\varphi \times H(\langle y, x \rangle_{C_c(H, A)}))g)^* f$$

for $x, y \in C_c(G, A) \subset X_H^G(A)$ and $g, f \in C_c(H, C)$. Thus Ψ converts the inner product on the internal tensor product $X_H^G(A) \otimes_{A \times H} (C \times H)$ to the usual one on $C \times_\epsilon H$.

To see that Ψ has dense range, and hence extends to an isomorphism of right-Hilbert bimodules, it is enough to approximate elements in $C_c(G, C)$ of the form $y \cdot f$, where $y \in C_c(G, C)$ and $f \in C_c(H, C)$, with elements in the range of Ψ . Moreover, a partition of unity argument shows that functions of the form $t \mapsto \varphi(x(t))\epsilon_t(c)$ for $x \in C_c(G, A)$ and $c \in C$ span an inductive-limit dense — and hence norm-dense — subspace of $C_c(G, C)$, so we can assume that y has this form. But now a routine calculation shows that if $g(h) = cf(h)$, then $\Psi(x \otimes g) = y \cdot f$ in $C_c(G, C)$; this implies that the range of Ψ is dense.

Another calculation using the analogue of (4.5) for the homomorphism $\varphi \times G$ shows that this isomorphism respects the left action of $A \times_\alpha G$. \square

To prove the commutativity of (4.4), we use a device from [ER96, §4]. Recall that if ${}_A X_B$ is an imprimitivity bimodule, then the linking algebra is the collection of 2×2 matrices

$$L(X) := \left\{ \begin{pmatrix} a & x \\ \tilde{z} & b \end{pmatrix} : a \in A, b \in B, x, z \in X \right\}.$$

with multiplication and involution given by

$$\begin{pmatrix} a & x \\ \tilde{z} & b \end{pmatrix} \begin{pmatrix} a' & x' \\ \tilde{z}' & b' \end{pmatrix} := \begin{pmatrix} aa' + {}_A \langle x, z' \rangle & a \cdot x' + x \cdot b' \\ (z \cdot a' + b \cdot z')^\sim & \langle z, x' \rangle_B + bb' \end{pmatrix}$$

and

$$\begin{pmatrix} a & x \\ \tilde{z} & b \end{pmatrix}^* := \begin{pmatrix} a^* & z \\ \tilde{x} & b^* \end{pmatrix}.$$

This has a unique complete C^* -norm, obtained, for example, by identifying $L(X)$ with the C^* -algebra $\mathcal{K}(X \oplus B)$ of compact operators on the Hilbert module direct sum $(X \oplus B)_B$ (see [RW98, page 50]). The matrices

$$p = p_{L(X)} := \begin{pmatrix} 1_{M(A)} & 0 \\ 0 & 0 \end{pmatrix} \quad \text{and} \quad q = q_{L(X)} := \begin{pmatrix} 0 & 0 \\ 0 & 1_{M(B)} \end{pmatrix}$$

define full projections in $M(L(X))$ which allow us to identify A, B and X with corners in $L(X)$; we can then use the projections p, q to break up a Hilbert $L(X)$ -module into modules over A and B . Our key technical result describes some relations among these submodules. It is a mild generalization of [ER96, Lemma 4.6].

Proposition 4.3. *Suppose ${}_A X_B$ and ${}_C Y_D$ are imprimitivity bimodules, and Z is a right-Hilbert $L(X)$ – $L(Y)$ bimodule. Then*

- (i) $pZp = p_{L(X)} Z p_{L(Y)}$ is a right-Hilbert A – C bimodule;
- (ii) $qZq = q_{L(X)} Z q_{L(Y)}$ is a right-Hilbert B – D bimodule;
- (iii) $pZq = p_{L(X)} Z q_{L(Y)}$ is a right-Hilbert A – D bimodule;

- (iv) *there is an isomorphism $\Phi : X \otimes_B qZq \rightarrow pZq$ of right-Hilbert A - D modules such that $\Phi(x \otimes z) = x \cdot z$;*
- (v) *there is an isomorphism $\Psi : pZp \otimes_C Y \rightarrow pZq$ of right-Hilbert A - D modules such that $\Phi(w \otimes y) = w \cdot y$.*

Remark 4.4. Together, the last two parts of Proposition 4.3 say that the diagram

$$\begin{array}{ccc}
 A & \xrightarrow{pZp} & C \\
 \downarrow X & \searrow pZq & \downarrow Y \\
 B & \xrightarrow{qZq} & D
 \end{array}$$

commutes in \mathcal{C} . If we had such a proposition when X and Y are just right-Hilbert bimodules, and the linking algebras are by definition $L(\kappa(X)X_B)$ and $L(\kappa(Y)Y_D)$, we could avoid having to factor our morphisms. However, we need to know that X and Y are imprimitivity bimodules to identify the corners $pL(X)p$ and $pL(Y)p$ with A and C . Since C in particular occurs in the middle of the internal tensor products in (v), it is hard to see how this hypothesis might be avoided.

Lemma 4.5. *Suppose ${}_EW_F$ is a right-Hilbert bimodule and $P \in M(E)$, $Q \in M(F)$ are full projections. Then PWQ is a right-Hilbert PEP - QFQ bimodule.*

Proof. Since PWQ is certainly a PEP - QFQ submodule, and since

$$\langle PWQ, PWQ \rangle_F = Q \langle PW, PW \rangle_F Q \subset QFQ,$$

we only have to check nondegeneracy of the left action and fullness on the right. For nondegeneracy, we use the fullness of P to see that $PEP \cdot WQ = PEP \cdot EWQ = P(EPE) \cdot WQ$ is dense in PWQ . For fullness, we use the fullness of P again to see that

$$\begin{aligned}
 \langle PWQ, PWQ \rangle_F &= Q \langle PE \cdot W, PE \cdot W \rangle_F Q \\
 &= Q \langle W, EP \cdot PE \cdot W \rangle_F Q = Q \langle W, E \cdot W \rangle_F Q
 \end{aligned}$$

is dense in QFQ . □

Proof of Proposition 4.3. For (i), apply Lemma 4.5 with $P = p_{L(X)} \in M(L(X))$ and $Q = p_{L(Y)} \in M(L(Y))$, and note that $A = PL(X)P$ and $C = QL(Y)Q$ because X and Y are imprimitivity bimodules. Parts (ii) and (iii) follow similarly.

For part (iv), we first note that because $(x, z) \mapsto x \cdot z$ is bilinear, there is a well-defined map Φ on the algebraic tensor product $X \odot qZq$ with the required property. We next verify that Φ preserves the inner product: if $x, x' \in X$ and $z, z' \in qZq$, then the inner product $\langle x, x' \rangle_B$ is given in $L(X)$ by $\langle x, x' \rangle_B = x^* x'$; more formally,

$$\begin{pmatrix} 0 & 0 \\ 0 & \langle x, x' \rangle_B \end{pmatrix} = \begin{pmatrix} 0 & x \\ 0 & 0 \end{pmatrix}^* \begin{pmatrix} 0 & x' \\ 0 & 0 \end{pmatrix}.$$

Thus

$$\begin{aligned}
 \langle x \otimes z, x' \otimes z' \rangle_D &= \langle z, \langle x, x' \rangle_B \cdot z' \rangle_D \\
 &= \langle z, (x^* x') \cdot z' \rangle_D \\
 &= \langle x \cdot z, x' \cdot z' \rangle_D \\
 &= \langle \Phi(x \otimes z), \Phi(x' \otimes z') \rangle_D,
 \end{aligned}$$

and Φ extends to an isometry of $X \otimes_B qZq$ into pZq . To see that Φ has dense range and is therefore onto, note that $L(X)$ acts nondegenerately on Z , so

$$\text{range } \Phi \supset pL(X)q \cdot qZq = pL(X)q \cdot q(L(X) \cdot Z)q = p(L(X)qL(X)) \cdot Zq$$

is dense because q is full.

For (v), note that the $D = qL(Y)q$ -valued inner product is given by the product in $L(Y)$, so

$$\begin{aligned}
 \langle w \otimes y, w' \otimes y' \rangle_D &= \langle y, \langle w, w' \rangle_C \cdot y' \rangle_D \\
 &= y^* \langle w, w' \rangle_{L(Y)} y' \\
 &= \langle w \cdot y, w' \cdot y' \rangle_{L(Y)},
 \end{aligned}$$

which is $\langle w \cdot y, w' \cdot y' \rangle_D$ because $w \cdot y$ and $w' \otimes y'$ belong to pZq . Thus $w \otimes y \mapsto w \cdot y$ extends to an isometry Ψ , as claimed, and Ψ is surjective because p is full and $L(Y)$ acts nondegenerately on Z . \square

Back to the proof of Theorem 4.1. Consider the action $\delta = \begin{pmatrix} \epsilon & \eta \\ \tilde{\eta} & \beta \end{pmatrix}$ of G on $L(Y)$; by Green's theorem, $Z = X_H^G(L(Y))$ is then a right-Hilbert $(L(Y) \times_\delta G) - (L(Y) \times_\delta H)$ bimodule. To show that (4.4) commutes, we aim to apply Proposition 4.3 to Z , but this requires that we first identify $L(Y) \times_\delta G$ with $L(Y \times_\eta G)$, and similarly for H . Now the dense subalgebra $C_c(G, L(Y))$ consists of 2×2 matrices with entries in $C_c(G, C)$, $C_c(G, Y)$, etc., and the diagonal corners have their usual $*$ -algebraic structures as subalgebras of $C \times_\epsilon G$ and $B \times_\beta G$. The norms are the same too: every covariant representation of $(L(Y), \delta)$ restricts to a covariant representation of (B, β) , and the inducing construction of Example 2.8 shows that every covariant representation of (B, β) extends to a covariant representation of $(L(Y), \delta)$. So we have embeddings of $C \times_\epsilon G$ and $B \times_\beta G$ as corners in $L(Y) \times_\delta G$. The bimodule crossed product $Y \times_\eta G$ also embeds: because $B \times G$ embeds isometrically, the norm of a function

$$t \mapsto \begin{pmatrix} 0 & f(t) \\ 0 & 0 \end{pmatrix} \in C_c(G, L(Y)) \subset L(Y) \times G$$

is just $\|\langle f, f \rangle_{B \times G}\|$. These embeddings combine to give an isomorphism of $L(Y \times_\eta G)$ onto $L(Y) \times_\delta G$, as desired. (This observation is not new: in [Com84], Combes *defines* $Y \times_\eta G$ to be the corner in $L(Y) \times_\delta G$.) In exactly the same way, $L(Y \times_\eta H) \cong L(Y) \times_\delta H$.

We next have to identify the corners $pX_H^G(L(Y), \delta)p$ and $qX_H^G(L(Y), \delta)q$ with $X_H^G(C, \epsilon)$ and $X_H^G(B, \beta)$. But viewing elements of $C_c(G, L(Y)) \subset X_H^G(L(Y), \delta)$ as matrices of functions, and similarly for $C_c(G, L(Y)) \subset L(Y) \times_\delta G$ and $C_c(H, L(Y)) \subset L(Y) \times_\delta H$, gives the desired identifications. It now follows from Proposition 4.3 that

$$(Y \times_\eta G) \otimes_{B \times G} X_H^G(B, \beta) \cong pX_H^G(L(Y), \delta)q \cong X_H^G(C, \epsilon) \otimes_{C \times H} (Y \times_\eta H)$$

as right-Hilbert $(C \times_\epsilon G) - (B \times_\beta H)$ bimodules. In other words, (4.4) commutes in the category \mathcal{C} .

We have now established the first part of Theorem 4.1, the natural transformation between the functors $(A, \alpha) \mapsto A \times_\alpha G$ and $(A, \alpha) \mapsto A \times_\alpha H$. For the second part, we have to prove that (4.2) commutes. We follow the same procedure as before: factor a given morphism $[{}_A X_B, \gamma]$ as $[{}_C Y_B, \eta] \circ [\varphi, \epsilon]$, and prove that two separate diagrams commute. To see the commutativity of

$$\begin{array}{ccc} (A \otimes C_0(G/H)) \times_{\alpha \otimes \tau} G & \xrightarrow{X_H^G(A, \alpha)} & A \times_\alpha H \\ (\varphi \otimes \text{id}) \times G \downarrow & & \downarrow \varphi \times H \\ (C \otimes C_0(G/H)) \times_{\epsilon \otimes \tau} G & \xrightarrow{X_H^G(C, \epsilon)} & C \times_\epsilon H, \end{array}$$

we merely note that the isomorphism of Lemma 4.2 respects the left action of $C_0(G/H)$, and hence is also an isomorphism of right-Hilbert $((A \otimes C_0(G/H)) \times_{\alpha \otimes \tau} G) - (C \times_\epsilon H)$ bimodules. To see that

$$\begin{array}{ccc} (C \otimes C_0(G/H)) \times_{\epsilon \otimes \tau} G & \xrightarrow{X_H^G(C, \epsilon)} & C \times_\epsilon H \\ (Y \otimes C_0(G/H)) \times_{\eta \otimes \tau} G \downarrow & & \downarrow Y \times_\eta H \\ (B \otimes C_0(G/H)) \times_{\beta \otimes \tau} G & \xrightarrow{X_H^G(B, \beta)} & B \times_\beta H \end{array}$$

commutes, we again apply Proposition 4.3 to $Z = X_H^G(L(Y), \delta)$, this time viewed as a $L((Y \otimes C_0(G/H)) \times_{\eta \otimes \tau} G) - L(Y \times_\eta H)$ imprimitivity bimodule. To do so, we have to identify $L((Y \otimes C_0(G/H)) \times_{\eta \otimes \tau} G)$ with $(L(Y) \otimes C_0(G/H)) \times_{\delta \otimes \tau} G$, but there is no essential difference in the argument.

This concludes the proof of our main theorem. \square

Remark 4.6. As an example of how the present set of ideas can simplify things, notice that the commutativity of diagram (4.4) implies that for each covariant representation $\rho \times V$ of $B \times_\beta H$, the representation $(Y \times_\eta G) - \text{Ind}(\text{Ind}_H^G(\rho \times V))$ is equivalent to $\text{Ind}_H^G((Y \times_\eta H) - \text{Ind}(\rho \times V))$. This is the content of [Ech94, Theorem 3], or at least the content of its proof (the twists in that theorem go along for free by Remark (2) on page 174 of [Ech94] and [Gre78, Corollary 5]). The point is that the arguments we give here are more elegant as well as more powerful.

REFERENCES

- [CMW84] R. Curto, P. Muhly, and D. Williams. Cross products of strongly Morita equivalent C^* -algebras. *Proc. Amer. Math. Soc.*, 90:528–530, 1984.
- [Com84] F. Combes. Crossed products and Morita equivalence. *Proc. London Math. Soc.*, 49:289–306, 1984.
- [Con94] Alain Connes. *Noncommutative Geometry*. Academic Press, San Diego, 1994.
- [Ech94] S. Echterhoff. Morita equivalent twisted actions and a new version of the Packer-Raeburn stabilization trick. *J. London Math. Soc.*, 50:170–186, 1994.
- [ER95] S. Echterhoff and I. Raeburn. Multipliers of imprimitivity bimodules and Morita equivalence of crossed products. *Math. Scand.*, 76:289–309, 1995.
- [ER96] S. Echterhoff and I. Raeburn. The stabilisation trick for coactions. *J. reine angew. Math.*, 470:181–215, 1996.
- [Gre78] P. Green. The local structure of twisted covariance algebras. *Acta Math.*, 140:191–250, 1978.
- [Kas88] G. G. Kasparov. Equivariant KK -theory and the Novikov conjecture. *Invent. Math.*, 91:147–201, 1988.
- [Kas95] G. G. Kasparov. K -theory, group C^* -algebras, and higher signatures (conspectus). In *Novikov conjectures, index theorems, and rigidity, Vol 1 (Oberwolfach, 1993)*, volume 226 of *London Math. Soc. Lecture Note Ser.*, pages 101–146. Cambridge University Press, 1995.
- [KQR97] S. Kaliszewski, J. Quigg, and I. Raeburn. Duality of restriction and induction for C^* -coactions. *Trans. Amer. Math. Soc.*, 349:2085–2113, 1997.
- [Lan95] E. C. Lance. *Hilbert C^* -Modules*, volume 210 of *London Math. Soc. Lecture Note Ser.* Cambridge University Press, 1995.
- [Men64] E. Mendelson. *Introduction to Mathematical Logic*. Van Nostrand, 1964.
- [MS98] Paul Muhly and Baruch Solel. Tensor algebras over C^* -correspondences: representations, dilations, and C^* -envelopes. *J. Funct. Anal.*, 158:389–457, 1998.
- [Pim97] M. V. Pimsner. A class of C^* -algebras generalizing both Cuntz-Krieger algebras and crossed products by \mathbf{Z} . In D.-V. Voiculescu, editor, *Free Probability Theory*, volume 12 of *Fields Inst. Commun.*, pages 189–212. Amer. Math. Soc., 1997.
- [Rae88] I. Raeburn. On crossed products and Takai duality. *Proc. Edinburgh Math. Soc.*, 31:321–330, 1988.
- [RW98] I. Raeburn and D. P. Williams. *Morita equivalence and continuous-trace C^* -algebras*. American Mathematical Society, 1998.
- [Tak67] M. Takesaki. Covariant representations of C^* -algebras and their locally compact automorphism groups. *Acta. Math.*, 119:273–303, 1967.

FACHBEREICH MATHEMATIK UND INFORMATIK, UNIVERSITY OF MÜNSTER, 48149 MÜNSTER,
GERMANY

E-mail address: `echters@uni-muenster.de`

DEPARTMENT OF MATHEMATICS, ARIZONA STATE UNIVERSITY, TEMPE, ARIZONA 85287

E-mail address: `kaz@math.la.asu.edu`

DEPARTMENT OF MATHEMATICS, ARIZONA STATE UNIVERSITY, TEMPE, ARIZONA 85287

E-mail address: `quigg@math.la.asu.edu`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF NEWCASTLE, NEW SOUTH WALES 2308,
AUSTRALIA

E-mail address: `iaain@maths.newcastle.edu.au`